

Complete Summary

GUIDELINE TITLE

Counselling. In: Canadian consensus guidelines on human papillomavirus.

BIBLIOGRAPHIC SOURCE(S)

Steben M, Money DM. Counselling. In: Canadian consensus guidelines on human papillomavirus. J Obstet Gynaecol Can 2007 Aug;29(8 Suppl 3):S55-6. [6 references]

GUIDELINE STATUS

This is the current release of the guideline.

COMPLETE SUMMARY CONTENT

SCOPE
METHODOLOGY - including Rating Scheme and Cost Analysis
RECOMMENDATIONS
EVIDENCE SUPPORTING THE RECOMMENDATIONS
BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS
QUALIFYING STATEMENTS
IMPLEMENTATION OF THE GUIDELINE
INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT
CATEGORIES
IDENTIFYING INFORMATION AND AVAILABILITY
DISCLAIMER

SCOPE

DISEASE/CONDITION(S)

Human papillomavirus (HPV) and HPV related diseases

GUIDELINE CATEGORY

Counseling

CLINICAL SPECIALTY

Family Practice
Infectious Diseases
Internal Medicine
Obstetrics and Gynecology

Pediatrics
Preventive Medicine

INTENDED USERS

Advanced Practice Nurses
Allied Health Personnel
Nurses
Physician Assistants
Physicians
Public Health Departments

GUIDELINE OBJECTIVE(S)

To promote guidelines for health care providers on the key aspects of human papillomavirus (HPV) infection and the management of HPV-related disease in the new era of vaccine availability

TARGET POPULATION

Sexually active women and adolescent girls

INTERVENTIONS AND PRACTICES CONSIDERED

1. Use of the "5 R's of Counselling" (reach out, review, resource, reassure, reiterate)
2. Help patients participate in their care
3. Discussion of adolescent sexuality

MAJOR OUTCOMES CONSIDERED

Not stated

METHODOLOGY

METHODS USED TO COLLECT/SELECT EVIDENCE

Searches of Electronic Databases

DESCRIPTION OF METHODS USED TO COLLECT/SELECT THE EVIDENCE

Medline and Cochrane databases were searched for articles from January 1995 to March 2007 on subjects related to Human papillomavirus (HPV) infection, HPV vaccination, HPV-related disease, Pap testing, and specific consideration of management.

NUMBER OF SOURCE DOCUMENTS

Not stated

METHODS USED TO ASSESS THE QUALITY AND STRENGTH OF THE EVIDENCE

Weighting According to a Rating Scheme (Scheme Given)

RATING SCHEME FOR THE STRENGTH OF THE EVIDENCE

Quality of Evidence Assessment*

I: Evidence obtained from at least one properly randomized controlled trial.

II-1: Evidence obtained from well-designed controlled trials without randomization.

II-2: Evidence obtained from well-designed cohort (prospective or retrospective) or case-control studies, preferably from more than one center or research group.

II-3: Evidence obtained from comparison between times or places with or without the intervention. Dramatic results in uncontrolled experiments (such as the results of treatment with penicillin in the 1940s) could also be regarded as this type of evidence.

III: Opinions of respected authorities, based on clinical experience, descriptive studies, or reports of expert committees.

*The quality of evidence reported in these guidelines has been adapted from The Evaluation of Evidence criteria described in the Canadian Task Force on Preventive Health Care.

METHODS USED TO ANALYZE THE EVIDENCE

Systematic Review

DESCRIPTION OF THE METHODS USED TO ANALYZE THE EVIDENCE

All study types were reviewed. Randomized controlled trial results were considered evidence of the highest quality, followed by results of cohort studies. Key individual studies on which the recommendations are based are referenced. Supporting data for each recommendation were summarized with evaluative comments and references.

METHODS USED TO FORMULATE THE RECOMMENDATIONS

Expert Consensus

DESCRIPTION OF METHODS USED TO FORMULATE THE RECOMMENDATIONS

Not stated

RATING SCHEME FOR THE STRENGTH OF THE RECOMMENDATIONS

Classification of Recommendations* †

- A. There is good evidence to recommend the clinical preventive action
- B. There is fair evidence to recommend the clinical preventive action
- C. The existing evidence is conflicting and does not allow to make a recommendation for or against use of the clinical preventive action; however, other factors may influence decision-making
- D. There is fair evidence to recommend against the clinical preventive action
- E. There is good evidence to recommend against the clinical preventive action
- I. There is insufficient evidence (in quantity or quality) to make a recommendation; however, other factors may influence decision-making

*Woolf SH, Battista RN, Angerson GM, Logan AG, Eel W. Canadian Task Force on Preventive Health Care. New grades for recommendations from the Canadian Task Force on Preventive Health Care. Can Med Assoc J 2003;169(3):207-8.

† Recommendations included in these guidelines have been adapted from the Classification of Recommendations criteria described in the Canadian Task Force on Preventive Health Care.

COST ANALYSIS

A formal cost analysis is included in *Chapter 7: Cost-Benefit Analysis of HPV Vaccination* in the original guideline document.

METHOD OF GUIDELINE VALIDATION

Internal Peer Review

DESCRIPTION OF METHOD OF GUIDELINE VALIDATION

These guidelines were prepared by the human papillomavirus (HPV) Consensus Guidelines Committee and approved by the Executive and Council of the Society of Obstetricians and Gynaecologists of Canada.

RECOMMENDATIONS

MAJOR RECOMMENDATIONS

The grades of recommendations (A-E and I) and levels of evidence (I, II-1, II-2, II-3, and III) are defined at the end of the "Major Recommendations" field.

1. A diagnosis of human papillomavirus (HPV) infection or its complications results in a wide range of emotional responses. Physicians should assess the impact the diagnosis has had on the patient and help her work through the emotional responses. **IIIA**

2. Health care providers should proactively discuss issues of sexuality with their patients. **IIIA**

Definitions

Levels of Evidence*

I: Evidence obtained from at least one properly randomized controlled trial.

II-1: Evidence obtained from well-designed controlled trials without randomization.

II-2: Evidence obtained from well-designed cohort or case-control studies, preferably from more than one center or research group.

II-3: Evidence obtained from comparison between times or places with or without the intervention. Dramatic results in uncontrolled experiments (such as the results of treatment with penicillin in the 1940s) could also be regarded as this type of evidence.

III: Opinions of respected authorities, based on clinical experience, descriptive studies, or reports of expert committees.

*The quality of evidence reported in these guidelines has been adapted from The Evaluation of Evidence criteria described in the Canadian Task Force on Preventive Health Care.

Grades of Recommendations* †

A. There is good evidence to recommend the clinical preventive action

B. There is fair evidence to recommend the clinical preventive action

C. The existing evidence is conflicting and does not allow to make a recommendation for or against use of the clinical preventive action; however, other factors may influence decision-making

D. There is fair evidence to recommend against the clinical preventive action

E. There is good evidence to recommend against the clinical preventive action

I. There is insufficient evidence (in quantity or quality) to make a recommendation; however, other factors may influence decision-making

* Woolf SH, Battista RN, Angerson GM, Logan AG, Eel W. Canadian Task Force on Preventive Health Care. New grades for recommendations from the Canadian Task Force on Preventive Health Care. Can Med Assoc J 2003;169(3):207-8.

† Recommendations included in these guidelines have been adapted from the Classification of Recommendations criteria described in the Canadian Task Force on Preventive Health Care

CLINICAL ALGORITHM(S)

None provided

EVIDENCE SUPPORTING THE RECOMMENDATIONS

TYPE OF EVIDENCE SUPPORTING THE RECOMMENDATIONS

The type of supporting evidence is identified and graded for each recommendation (see "Major Recommendations").

BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS

POTENTIAL BENEFITS

- Appropriate management of patients with human papillomavirus (HPV) infection
- Timely crisis intervention and assistance with coping
- Promotion of self-care and shared decision-making
- Reduction in HPV transmission

POTENTIAL HARMS

Not stated

QUALIFYING STATEMENTS

QUALIFYING STATEMENTS

This document reflects emerging clinical and scientific advances on the date issued and is subject to change. The information should not be construed as dictating an exclusive course of treatment or procedure to be followed. Local institutions can dictate amendments to these opinions. They should be well documented if modified at the local level. None of these contents may be reproduced in any form without prior written permission of the Society of Obstetricians and Gynaecologists of Canada (SOGC).

IMPLEMENTATION OF THE GUIDELINE

DESCRIPTION OF IMPLEMENTATION STRATEGY

An implementation strategy was not provided.

INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT CATEGORIES

IOM CARE NEED

Getting Better
Staying Healthy

IOM DOMAIN

Effectiveness
Patient-centeredness
Timeliness

IDENTIFYING INFORMATION AND AVAILABILITY

BIBLIOGRAPHIC SOURCE(S)

Steben M, Money DM. Counselling. In: Canadian consensus guidelines on human papillomavirus. J Obstet Gynaecol Can 2007 Aug;29(8 Suppl 3):S55-6. [6 references]

ADAPTATION

Not applicable: The guideline was not adapted from another source.

DATE RELEASED

2007 Aug

GUIDELINE DEVELOPER(S)

Society of Obstetricians and Gynaecologists of Canada - Medical Specialty Society

SOURCE(S) OF FUNDING

The development of this consensus guideline was supported by unrestricted educational grants from Cytyc Canada, Digene Corporation, Graceway Canada, GlaxoSmithKline Inc., Merck Frosst Canada Ltd., and Roche Diagnostics Canada.

GUIDELINE COMMITTEE

HPV Consensus Guidelines Committee

COMPOSITION OF GROUP THAT AUTHORED THE GUIDELINE

Primary Authors: Marc Steben, MD, Montreal QC and Deborah M. Money, MD, FRCSC, Vancouver BC

Committee Members: Deborah M. Money, MD, FRCSC (*Chair*) Vancouver BC; Michel Roy, MD, FRCSC (*Chair*) Quebec City QC; Lisa Allen, MD, FRCSC, Toronto ON; Monica Brewer, MD, Quispamsis NB; Peter Bryson, MD, FRCSC, Kingston ON; Gerald Evans, MD, FRCSC, Kingston ON; Jean-Yves Frappier, MD, FRCSC, Montreal QC; Mary Anne Jamieson, MD, FRCSC, Kingston ON; Charles Lynde, MD, FRCPC, Markham ON; K. Joan Murphy, MD, FRCSC, Toronto ON; Diane M. Provencher, MD, FRCSC, Montreal QC; Michael Shier, MD, FRCSC, Toronto ON; Marc Steben, MD, Montreal QC; Robert Lerch, Public Health Agency of Canada,

Ottawa ON; André Lalonde, MD, FRCSC, Executive Vice-President, Ottawa ON;
Vyta Senikas, MD, FRCSC, Associate Executive Vice-President, Ottawa ON

FINANCIAL DISCLOSURES/CONFLICTS OF INTEREST

Disclosure statements have been received from all members of the committees.

ENDORSER(S)

Canadian Association for Adolescent Health - Medical Specialty Society
Canadian Pediatric and Adolescent Gynaecology and Obstetrics Committee -
Medical Specialty Society
Federation of Medical Women of Canada - Professional Association
Quebec Association of Pediatricians - State/Local Government Agency [Non-U.S.]
Society of Canadian Colposcopists - Professional Association
Society of Gynecologic Oncologists of Canada - Disease Specific Society

GUIDELINE STATUS

This is the current release of the guideline.

GUIDELINE AVAILABILITY

Electronic copies: Available in Portable Document Format (PDF) from the [Society of Obstetricians and Gynaecologists of Canada Web site](#).

Print copies: Available from the Society of Obstetricians and Gynaecologists of Canada, La société des obstétriciens et gynécologues du Canada (SOGC) 780 promenade Echo Drive Ottawa, ON K1S 5R7 (Canada); Phone: 1-800-561-2416

AVAILABILITY OF COMPANION DOCUMENTS

None available

PATIENT RESOURCES

None available

NGC STATUS

This NGC summary was completed by ECRI Institute on July 8, 2009. The information was verified by the guideline developer on July 14, 2009.

COPYRIGHT STATEMENT

This NGC summary is based on the original guideline, which is subject to the guideline developer's copyright restrictions.

DISCLAIMER

NGC DISCLAIMER

The National Guideline Clearinghouse™ (NGC) does not develop, produce, approve, or endorse the guidelines represented on this site.

All guidelines summarized by NGC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public or private organizations, other government agencies, health care organizations or plans, and similar entities.

Guidelines represented on the NGC Web site are submitted by guideline developers, and are screened solely to determine that they meet the NGC Inclusion Criteria which may be found at <http://www.guideline.gov/about/inclusion.aspx>.

NGC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or clinical efficacy or effectiveness of the clinical practice guidelines and related materials represented on this site. Moreover, the views and opinions of developers or authors of guidelines represented on this site do not necessarily state or reflect those of NGC, AHRQ, or its contractor ECRI Institute, and inclusion or hosting of guidelines in NGC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding guideline content are directed to contact the guideline developer.

[Copyright/Permission Requests](#)

Date Modified: 8/10/2009

